



1/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form A measured with variable slits.

**Fig. 1**


2/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form B measured with variable slits.

*Fig. 2*


3/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form C measured with variable slits.

**Fig. 3**


4/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form E measured

*Fig. 4*


5/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form F measured with variable slits.

**Fig. 5**


6/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form G measured with variable slits.

**Fig. 6**


7/9



An X-ray powder diffractogram of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form H measured with variable slits.

*Fig. 7*

8/9




A Raman spectrum of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazo[1,2-a]pyridine-6-carboxamide mesylate salt form A

**Fig. 8**



9/9



A Raman spectrum of 2,3-dimethyl-8-(2,6-dimethylbenzylamino)-N-hydroxyethyl-imidazol[1,2-a]pyridine-6-carboxamide mesylate salt form B

**Fig. 9**